

Lorentz and CPT violations in QED : a short comment on recent controversies.

Guy Bonneau*

Abstract

We rediscuss the recent controversy on a possible Chern-Simons like term generated through radiative corrections in QED with a CPT violating term. We emphasize the fact that any absence of an *a priori* divergence should be explained by some symmetry or some non-renormalisation theorem : otherwise, no prediction can be made on the corresponding quantity.

PACS codes : 11.10.Gh, 11.30.Er, 12.20.-m

Keywords : Ward identities, Radiative corrections, CPT violation, Chern-Simons

In the past years, the interesting issue of a possible spontaneous breaking of Lorentz invariance at low energy has been considered : this issue also led to CPT breaking [1, 2, 3]. In particular, the general Lorentz-violating extension of the minimal $SU(3) \times SU(2) \times U(1)$ standard model has been discussed : as many breaking terms are allowed, people look for possible constraints coming from experimental results as well as from renormalisability requirements and anomaly cancellation.

In that respect, there arose a controversy on a possible Chern-Simons like term generated through radiative corrections [2-16]. This phenomenon was studied in QED, an abelian gauge theory, as a part of the standard model. The aim of this note is to clarify the discrepancies and insit upon the need of an explanation when some one-loop divergence does not appear.

The Lagrangian density is :

$$\begin{aligned} \mathcal{L} &= \mathcal{L}_0 + \mathcal{L}_1 + \mathcal{L}_2 \\ a) \mathcal{L}_0 &= \bar{\psi}(i \not{\partial} - m - e \not{A})\psi - \frac{1}{4}F_{\mu\nu}^2 - \frac{1}{2\alpha}(\partial A)^2 + \frac{1}{2}\lambda^2 A_\mu^2 \\ &\quad \text{where } \alpha \text{ is the gauge parameter and } \lambda \text{ an infra - red regulator photon mass ,} \\ b) \mathcal{L}_1(x) &= -b^\mu \bar{\psi}(x)\gamma_\mu \gamma^5 \psi(x), \quad \text{where } b^\mu \text{ is a fixed vector,} \\ c) \mathcal{L}_2(x) &= \frac{1}{2}c^\mu \epsilon_{\mu\nu\rho\sigma} F^{\nu\rho}(x) A^\sigma(x), \quad \text{where } c^\mu \text{ is a fixed vector.} \end{aligned} \tag{1}$$

Other breakings could be considered (see a discussion in the first paper of [2]), but we simplify and require charge conjugation invariance, which selects $\mathcal{L}_1(x)$ and $\mathcal{L}_2(x)$. Note for further reference that experiments on the absence of birefringence of light in vacuum put very restrictive limits on the value of c^μ , typically for a timelike c^μ , $c^0/m \leq 10^{-38}$ [2].

- In order to avoid the difficulties resulting from the new poles in the propagators, as in [9], we take into account the smallness of the breakings and include them into the interaction Lagrangian density as super-renormalisable couplings (see in [2] a discussion on the alternative choice of a complete propagator). Moreover, we define the photon and electron masses by the same normalisation conditions as in ordinary Q.E.D., *e.g.*

$$\langle \psi(p) \bar{\psi}(-p) \rangle^{prop.}_{p^2=m, b=c=0} = 0, \quad \dots$$

According to standard results in renormalisation theory, these breakings add new terms in the primitively divergent proper Green functions. By power counting, these are

$$\Gamma_{\mu\nu}(p, -p), \quad \Sigma(p, -p), \quad \Gamma^\rho(p, q, -(p+q)) \text{ and } \Gamma_{\mu\nu\rho\sigma}(p_1, p_2, p_3, -(p_1+p_2+p_3)),$$

respectively the photon and electron 2-points proper Green functions, the photon-electron proper vertex function and the photon 4-point proper Green function. The corresponding overall divergences (sub-divergences being properly subtracted) are polynomial in the momenta and masses ¹:

$$\begin{aligned} \Gamma_{\mu\nu}(p, -p) |_{div} &= a_1[g_{\mu\nu}p^2 - p_\mu p_\nu] + a_2 p_\mu p_\nu + [a_3 m^2 + a_4 \lambda^2]g_{\mu\nu} + [a_5 b^\rho + a_6 c^\rho]\epsilon_{\mu\nu\rho\sigma} p^\sigma, \\ \Sigma(p, -p) |_{div} &= a_7 \not{p} + a_8 m + [a_9 b^\rho + a_{10} c^\rho]\gamma_\rho \gamma^5, \\ \Gamma^\rho(p, q, -(p+q)) |_{div} &= a_{11} \gamma^\rho. \\ \Gamma_{\mu\nu\rho\sigma}(p_i) |_{div} &= a_{12}[g_{\mu\nu}g_{\rho\sigma} + g_{\mu\rho}g_{\nu\sigma} + g_{\mu\sigma}g_{\rho\nu}]. \end{aligned} \tag{2}$$

¹ C invariance has been used. The Ward identity (4) will relate some of these parameters : $a_2 = a_3 = a_4 = 0$, $a_{11} = ea_7$, $a_{12} = 0$.

All parameters $a_i[e, \frac{b^2}{m^2}, \frac{c^2}{m^2}, \frac{b \cdot c}{m^2}]$ (positions and residues of the poles in propagators, couplings at zero momenta,..) - but for the unphysical, non renormalised ones (as the longitudinal photon propagator (gauge parameter α) and photon regulator mass λ^2 for unbroken QED) - require normalisation conditions, a point which has often been missed since the successes of **minimal** dimensionnal regularisation scheme [17] but is stressed in the recent reviews [10, 11] and in a computation on QED at finite temperature [16]. In particular we shall require 2 new normalisation conditions to fix the breaking parameters b^μ and c^ν :

$$\begin{aligned} b^\mu &= -\frac{i}{4} \text{Tr}[\gamma^\mu \gamma^5 < \psi(p) \bar{\psi}(-p) >^{prop.}]|_{p=0}, \\ c^\mu &= \frac{1}{12} \epsilon^{\mu\nu\rho\sigma} \frac{\partial}{\partial p^\sigma} < A_\nu(p) A_\rho(-p) >^{prop.}|_{p=0}. \end{aligned} \quad (3)$$

Note that, contrary to $\mathcal{L}_1(x)$, the $\mathcal{L}_2(x)$ term also breaks the local gauge invariance of the Lagrangian density, but we emphasize the fact that - except for the unphysical part $\int [-\frac{1}{2\alpha}(\partial A)^2 + \frac{1}{2}\lambda^2 A_\mu^2]$ - the action $\Gamma = \int \mathcal{L}$ is invariant under local gauge transformations.

Then a Ward identity may be written :

$$\begin{aligned} &\int d^4x \left\{ \frac{1}{e} \partial_\mu \Lambda(x) \frac{\delta \Gamma}{\delta A_\mu(x)} + i \Lambda(x) [\bar{\psi}(x) \frac{\overrightarrow{\delta} \Gamma}{\delta \bar{\psi}(x)} - \frac{\Gamma \overleftarrow{\delta}}{\delta \psi(x)} \psi(x)] \right\} = \\ &= \int d^4x \left\{ -\frac{1}{e\alpha} \partial_\mu A^\mu(x) \square \Lambda(x) + \frac{\lambda^2}{e} A^\mu(x) \partial_\mu \Lambda(x) + \frac{1}{2e} \epsilon_{\alpha\beta\delta\mu} c^\alpha F^{\beta\delta}(x) \partial^\mu \Lambda(x) \right\} \\ &\Rightarrow W_x \Gamma \equiv \partial_\mu \frac{\delta \Gamma}{\delta A_\mu(x)} - ie [\bar{\psi}(x) \frac{\overrightarrow{\delta} \Gamma}{\delta \bar{\psi}(x)} - \frac{\Gamma \overleftarrow{\delta}}{\delta \psi(x)} \psi(x)] = \frac{1}{\alpha} [\square + \alpha \lambda^2] \partial_\mu A^\mu(x). \end{aligned} \quad (4)$$

Note that this equation is exactly the same as the one for ordinary QED.

As soon as we use a regularisation that respects the symmetries (gauge, Lorentz covariance and charge conjugation invariance), the perturbative proof of renormalisability reduces to the check that the $\mathcal{O}(\hbar)$ quantum corrections to the classical action $\Gamma : \Gamma_1 = \Gamma|_{class.} + \hbar \Delta$, constrained by the Ward identity (4) may be reabsorbed into the classical action through suitable renormalisations of the fields and parameters of the theory. This has been proved in [9].

There, some local sources have been introduced to define the **local** operators $\mathcal{L}_1(x)$ and $\mathcal{L}_2(x)$. Although this is only a technical tool, it has been criticised ² so we now discuss whether the quantization of $\Gamma = \int \mathcal{L}$, without a **local** definition of the operators $\mathcal{L}_1(x)$ and $\mathcal{L}_2(x)$, is possible.

- Of course, in ordinary QED, the axial current, being uncoupled, is absent from the Lagrangian density and so does not need to be defined as a quantum operator ; no axial vertex being present, *a fortiori* there is no axial anomaly and no triangle graph to consider.

² For example in page 3 of [11] : “Bonneau introduced external source fields for the axial vector current and the CS term, so the Ward identities he derived actually impose gauge invariance on Lagrangian density ...” This assertion is wrong as the Lagrangian density is not gauge invariant (moreover it has been gauge-fixed..) but, as proven in our analysis [9], the breaking of local gauge invariance is a soft one and may be seen as a complementary part in the gauge fixing.

On the contrary, in CPT-broken QED (1), new axial insertions enter the game, but they are integrated ones $\int d^4x \mathcal{L}_1(x)$ and $\int d^4x \mathcal{L}_2(x)$. Then, most of the authors in that subject argue that introducing local sources for the Lagrangian density breakings means adding supplementary conditions on the theory.

I have two answers, or rather two objections :

- During my studies at University, I was not taught - nor have I taught ! - how it is possible to **define** the **space-time integration** of some local quantity if this quantity itself is not defined at (nearly) every point of the space time (notice that here the involved quantity is a product of quantum fields at the same point of the space-time). On the contrary, space-time integration, *i.e.* vanishing incoming momentum, sometimes introduces new difficulties (IR divergences,...) ! And, to define a quantum local operator with the generating functional approach, I know no other way than the introduction of local sources, so it is what I did in [9].
- Leaving aside this mathematical objection, suppose that one has only the Ward identity (4) at hand to constrain the possible ultra-violet divergences (2). This is not sufficient to prove that the breakings introduce no new infinities : in particular, the Chern Simons term is of the right canonical dimension and quantum numbers and satisfies (4) :

$$p^\mu \Gamma_{\mu\nu}(p, -p) = 0 \quad \text{in particular} \Rightarrow p^\mu < [\int d^4x \mathcal{L}_1(x)] A^\mu(p) A^\nu(-p) > = 0.$$

So, first, we have no explanation of the fact that **all one-loop calculations of the CS contribution to the photon self-energy give a finite result** ($a_5 = a_6 = 0$), second, being unconstrained, its finite part (renormalised value) has to be fixed by a normalisation condition (a different situation than a radiative correction such as the (g-2) or the Lamb-shift for example). So, no prediction is possible and its value remains arbitrary, which is rather unsatisfactory.

I am really surprised that in the twenty or so papers devoted to that subject, I could not find one line of argument to explain this “experimental ” one-loop³ finiteness, except in the recent review by Pérez-Victoria [10] where the finiteness of the CS term is related to the one of the standard triangle graph in ordinary QED : however, in QED too, the finiteness of such a graph results from the gauge Ward identity on the **unintegrated axial vertex** 3-point function :

$$p_\nu < [\bar{\psi} \gamma^\mu \gamma^5 \psi](-p-q) A^\nu(p) A^\rho(q) > = q_\rho < [\bar{\psi} \gamma^\mu \gamma^5 \psi](-p-q) A^\nu(p) A^\rho(q) > = 0. \quad (5)$$

So, as long as no answer to these two objections has been given, either you consider an ill-defined and no predictive theory of broken-QED⁴ or you agree to consider an action in which every Lagrangian monomial is well-defined as a quantum operator.

Then in [9], I have proved that, being linear in the quantum field, the variation of $\mathcal{L}_2(x)$ in a local gauge transformation is soft : no essential difference occurs between local gauge invariance of the action and the “softly” broken local gauge invariance of the Lagrangian density. As a consequence, the theory (1) is consistent (even with no $\mathcal{L}_2(x)$ term) and the

³ If this finiteness was an “accidental” one, it would have no reason to hold at higher-loop order !

⁴ Some authors rightly conclude that the corresponding one-loop finite contribution is ambiguous [4, 8, 10, 11].

CS term has been shown to be unrenormalised, **to all orders of perturbation theory**. So, its experimental “vanishing” offers no constraint on the other CPT breaking term $\mathcal{L}_1(x)$.

- As a complement to the recent reviews [10, 11], let us now comment upon some points given in the literature :
 - Jackiw and Kosteletzky [5] never introduce any regulator. Then some of their relations are “delicate ones” : see for example for a divergent integral (after equ.12), the commutation of a derivation with respect to external momentum and the integration ⁵. If the integral in their equation (11) is computed with dimensional regularisation, a result $\left[\frac{\theta}{\sin \theta} - 1 \right]$ is found, and not simply $\frac{\theta}{\sin \theta}$ (with $p^2 = 4m^2 \sin^2 \theta/2$).
 - Moreover, in the absence of normalisation conditions or Ward identities fixing some ambiguities, the difference of two equivalent linearly divergent integrals gives an ambiguous logarithmic divergent one. Even when one uses a symmetric integration that suppresses the linearly (and eventually the logarithmically) divergent part, the finite part remains ambiguous. The “surface term” that comes from a shift in the integration momentum in a linearly divergent integral is a regulator dependent quantity : if one mimics the calculation in the appendix A5-2 of Jauch and Rohrlich standard book [18] with the dimensional scheme, one easily checks that no “surface term” occurs after a shift of the integration momenta [19]). Recall that this possibility of shifting internal momenta is needed to preserve gauge invariance in loop calculations (see for example [20, subsect.17.9]).
 - In a recent work [12], the one-loop calculation of the CS correction is done with the heat-kernel expansion and the Schwinger proper-time method, leading to a new finite result, claimed to be unambiguously determined. However,
 - * here again there is no explanation of the absence of infinities in the result : then the finite part is *a priori* ambiguous ⁶,
 - * other computations with the Schwinger proper-time method exist [13] and give a different result, proving at least that some “ambiguity” remains,
 - * some terms are lacking in this calculation : in particular a logarithmically divergent contribution to the CS term results from a thorough computation of the quantity given in equation (21) of [12] (in the absence of any precise criteria to substract infinite parts, this should not be a surprise).
 - It is difficult to see the difference often advocated between a first order (in b^μ) perturbative calculation and what is claimed to be a “non-perturbative unambiguous value”, but is, as a matter of fact, obtained with exactly the same standard triangle integrals [4]. Moreover, in [8] the computation is also done to all orders in the breaking parameter b^μ and it is explicitly verified that higher orders do not contribute to a possible correction to c^μ .

⁵ I cannot agree, and probably no teacher can agree, with the comment on this work given in Chen’s review ([11] p. 3): “*They ingeneously manipulated the linear divergent term in the loop integration ..*”. One should expect that such ideas will disappear in published works.

⁶ Remember that in Fujikawa’s calculation of the axial anomaly, gauge invariance was implemented through the basis used to compute the fermionic Jacobian : he chose eigenvectors of the operator $i \not{D} - e \not{A}$; another choice would allow the transfer of the axial anomaly to some vector anomaly (see also the discussion on the “minimal anomaly” in non-abelian gauge theory) [21].

However, in the first paper in [2], Colladay and Kosteleck'y gave a direct analysis of the **complete** classical fermion Green function as defined by $\mathcal{L}_0 + \mathcal{L}_1$. In particular they check that the anticommutator of two fermionic fields vanishes for space-like separations, in agreement with microcausality (at least for a time-like breaking b_μ). This confirms our analysis on the correctness of a theory with no classical CS term. Then, Adam and Klinkhamer show that the addition of a (radiatively generated) CS term $\mathcal{L}_2(x)$ with a time-like c^μ breaks microcausality [14]. As our non-renormalisation theorem ensures that, if absent at the classical level, the CS term will not appear in higher-loop order, microcausality will not be destroyed in higher-loop order.

Note also that the Lagrangian density $\mathcal{L}_0 + \mathcal{L}_2$ would not lead to a coherent theory as an (infinite) counterterm $\mathcal{L}_1$ appears at the one-loop order [9].

To summarize, we have proven that :

- The local gauge invariance of the Lagrangian density is destroyed by a $\mathcal{L}_2$ term (plus of course by the usual gauge fixing term) : but, being bilinear in the gauge field, $\mathcal{L}_2(x)$ behaves as a minor modification of the **gauge fixing term** as $\partial_\nu A^\nu$ remains a free field. As part of the “gauge term”, this $\mathcal{L}_2(x)$ is, as usual, not renormalised : so its all-order value is equal to its (arbitrarily chosen) classical one.
- A theory with a vanishing tree level Chern-Simons like breaking term is consistent as soon as it is correctly defined : thanks to the gauge invariance of the action, we have proven that the normalisation condition $c^\mu = 0$ may be enforced to all orders of perturbation theory.
- The 2-photon Green function receives definite (as they are finite by power counting) radiative corrections [9]

$$\simeq \frac{\hbar e^2}{12\pi^2} \frac{p^2}{m^2} \epsilon_{\mu\nu\rho\sigma} p^\sigma b^\rho + \dots$$

Recall the case of the electric charge : physically measurable quantities occur only through the p^2 dependence of the photon self-energy (as the Lamb-shift is a measurable consequence of a non-measurable charge renormalisation). Unfortunately, as Coleman and Glashow explained, the absence of birefringence of light in vacuum, *i.e.* the vanishing of the parameter c^μ , gives no constraint on the value of the other one b^μ . However, in [15, 16], CPT breaking in QED is studied at finite temperature : as is clearly emphasized in [16], the T dependance being independant on the normalisation condition at $T=0$, as soon as it is carefully computed it offers an unambiguous induced Chern-Simons like term at finite T and can give some information on b^μ .

Thanks : I have truly appreciated some correspondance with M. Pérez-Victoria.

References

- [1] S. Caroll, G. Field and R. Jackiw, *Phys. Rev.* **D 41** (1990) 1231.
- [2] D. Colladay and V. A. Kostelecky, *Phys. Rev.* **D55** (1997) 6760 ; *Phys. Rev.* **D58** (1998) 116002, and references therein.

- [3] S. Coleman and S. L. Glashow, *Phys. Rev.* **D59** (1999) 116008.
- [4] R. Jackiw, “When radiative corrections are finite but undetermined”, [hep-th/9903044].
- [5] R. Jackiw and V. A. Kostelecky, *Phys. Rev. Lett.* **82** (1999) 3572.
- [6] J.-M. Chung and P. Oh, *Phys. Rev.* **D60** (1999) 067702.
- [7] W. F. Chen, *Phys. Rev.* **D60** (1999) 085007.
- [8] J. M. Chung, *Phys. Lett.* **B461** (1999) 138 ; M. Pérez-Victoria, *Phys. Rev. Lett.* **83** (1999) 2518.
- [9] G. Bonneau, *Nucl. Phys.* **B593** (2001) 398, [hep-th/0008210].
- [10] M. Pérez-Victoria, *J. H. E. P.* **0104** (2001) 032, [hep-th/0102021].
- [11] W. F. Chen, “Issues on radiatively induced Lorentz and CPT violation in quantum electrodynamics”, [hep-th/0106035].
- [12] Yu. A. Sitenko, *Phys. Lett.* **B515** (2001) 414, [hep-th/0103215].
- [13] M. Chaichian, W. F. Chen and R. González Felipe, *Phys. Lett.* **B503** (2001) 215, [hep-th/0010129] ; J.-M. Chung and B. K. Chung, *Phys. Rev.* **D63** (2001) 105015, [hep-th/0101097].
- [14] C. Adam and F.R. Klinkhamer, *Phys. Lett.* **B513** (2001) 245, [hep-th/0105037].
- [15] J. R. S. Nascimento, R. F. Ribeiro and N. F. Svaiter, “Radiatively induced Lorentz and CPT violation in QED at finite temperature”, [hep-th/0012039].
- [16] L. Cervi, L. Griguolo and D. Seminara, *Phys. Rev.* **D64** (2001) 105003, [hep-th/0104022].
- [17] G. Bonneau, *Int. J. of Mod. Phys.* **A 5** (1990) 3831.
- [18] J. M. Jauch and F. Rohrlich, “The theory of Photons and Electrons”, Addison-Wesley Pub. Cie. (Reading, Ma., 1959).
- [19] K. J. Wilson, *Phys. Rev.* **D7** (1973) 2911, appendix.
- [20] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Fields*, McGraw-Hill Book Company.
- [21] K. Fujikawa, *Phys. Rev. Lett.* **42** (1979) 1195 ; *Phys. Rev.* **D21** (1980) 2848.